Instructions for completing the Prescription and Patient Support Program Enrollment Form

Adempas® riociguat tablets
0.5mg | 1mg | 1.5mg | 2mg | 2.5mg

Adempas Prescription (complete steps 1-3)

- Complete patient information and prescription details
- Provide medical and prescription insurance information
- AIM Coordination Center will complete benefit investigation to verify benefits and determine coverage

Note: Patients will be assessed for all available programs. Eligible patients will be automatically enrolled in the Adempas Co-pay Program

SECTION 1 Contact Information

By providing an email or phone, the patient agrees to receive automated calls, texts and/or email messages from myAIM Program administrator and the patient's pharmacy (CVS Specialty or Accredo) about your prescription and healthcare. Consent may be revoked at any time and is not a condition of the service. Carrier message and data rates may apply.

Patient Contact Information (* indicates required field)

Patient First Name*:	Patient Last Name*:			Birthdate* (MM/DD/YYYY):	Sex*: ☐Male ☐Female
Address*:	City*:	State*:	Zip Code*:	Preferred Phone*:	*OK to leave detailed message?* ☐Yes ☐No
Email: @	Preferred Language: ☐ English ☐ Spanish ☐ Other (specify) _____				
Alternate Contact Name:	Alternate Contact Phone:			Relationship to Patient:	

Prescriber Contact Information (* indicates required field)

Prescriber First Name*:	Prescriber Last Name*:			NPI*:
Address Line 1*:	Address Line 2:	City:	State:	Zip Code:
Office Contact:	Phone:	Fax:		

SECTION 2 Patient Information

Patient Information (* indicates required field)

Is Patient starting Adempas in a hospital setting? ☐**Yes** ☐**No** Start Date:_____ Discharge Date:_____

Does the patient have prescription coverage*? ☐Yes ☐No

Patient's local pharmacy:_____ Phone: _____

***PROVIDE ALL PATIENT INSURANCE INFORMATION, INCLUDING DRUG BENEFITS (FRONT AND BACK OF CARD) WITH THIS FORM.**

Please check one ICD-10 Code*:

Pulmonary arterial hypertension
☐ I27.0
☐ I27.21

Chronic thromboembolic pulmonary hypertension
☐ I27.24
☐ Inoperable
☐ Persistent/Recurrent

☐**OTHER (please specify)** _____

Therapy Status:
☐ Initial therapy (monotherapy or in combination)
☐ Add-on therapy
☐ Transition from other therapy

Prescriber will comply with all Surescripts†† terms and conditions including confidentiality, commercial messaging, privacy and security, applicable laws, and use of data. All Surescripts' disclaimers apply. A full list of terms and conditions is available at *https://ubc.com/surescriptsterms/*

SECTION 3 Prescription

Prescription (* indicates required field)
Note: NY Prescribers please submit prescription on an original NY State prescriptions blank. For all other States, send on a State-specific prescription blank if applicable for your State.

Starting dose*:	Titration schedule:
☐ **Adempas 1 mg tablet by mouth three times a day** ☐ Adempas 0.5 mg tablet by mouth three times a day Quantity: ☐ 30 day supply Refills:_____ Deliver to: ☐Patient Home ☐Prescriber Office	**Please check box for all dosages to be incorporated**: ☐ Based on patient's response per clinical evaluation of the physician or the nurse in consultation with the physician, the pharmacy is to provide the Adempas strength to accommodate titration needs of therapy. Adempas Tablets: 0.5 mg, 1 mg, 1.5 mg, 2 mg, 2.5 mg Directions: If systolic blood pressure is >95 mmHg and there are no signs/symptoms of hypotension, up titrate by 0.5 mg 3 times per day at intervals no sooner than 2 weeks to the highest tolerated dosage up to a maximum of 2.5 mg 3 times per day. If at any time, the patient has symptoms of hypotension, decrease the dosage by 0.5 mg 3 times daily. The established individual dose should be maintained. **Other special instructions:** _____ Titration Quantity: ☐ 30 day supply ☐ 14 day supply Refills: _____

I certify that the above information provided is accurate to the best of my knowledge. I appoint the Adempas AIM Program, on my behalf, to convey this prescription to the dispensing pharmacy. I understand that I may not delegate signature authority. Prescriber authorizes UBC to use the Surescripts Network† on Prescriber's behalf in connection with this prescription.

PRESCRIBER SIGNATURE REQUIRED	**Dispense as Written*:**	**Date*:**
	Substitutions Permitted*:	**Date*:**

Return this form along with patient insurance information to the Adempas Program via fax to 1-855-662-5200

†Surescripts is a consortium owned by some of the country's largest PBMs that offers information and technology services that support the electronic transmission of prescriptions between HCPs and other healthcare organizations.

Patient Support Enrollment (complete step 4)

SECTION 4 Patient Support Program Enrollment

Patient Support Program Enrollment

Bayer offers patient support services for Adempas patients that include: (A) nurses to support you in starting therapy and achieving your optimal dose, (B) insurance benefit verification for Adempas and financial assistance for eligible patients and (C) education about CTEPH and/or PAH as well as helpful tips for managing your Adempas therapy ("myAIM"). These Programs are entirely optional and you may enroll in one or all of these Programs. To enroll in myAIM, you will need to sign a HIPAA authorization in order for your healthcare provider and/or pharmacy to share your protected healthcare information with Bayer and the myAIM Program administrator.

You will remain enrolled in each Program that you select unless you opt-out by contacting myAIM via telephone at 1-855-423-3672 or until your HIPAA Authorization expires.

Please enroll me in: (check all that apply) ☐ A: Nursing ☐ B: Benefits Verification and Financial Assistance

☐ C: Digital Educational Information – e-mail address: _____@_____

Patient – please initial here to confirm your optional selections: _____

Patient can opt-out of any one of the above programs (or all) by contacting the AIM program.

Please also read, sign, and date the PATIENT HIPAA AUTHORIZATION at the end of this form.

PATIENT TO SIGN AND DATE

Patient Name (print): _____

Patient (or legal guardian) Signature*: _____ **Date (mm/dd/yyyy):** _____

If signed by a legal representative —
Print Name: _____ **Relationship to patient:** _____

At least 1 phone number is required

Prescribers in NY must submit prescriptions on NYS official Rx form, together with this form

Missing signatures and date **WILL** cause a delay in processing

Selections for patient programs and initial to confirm

Missing signatures and date **WILL** cause a delay in processing; printed name also required for processing

Adempas® (riociguat) Prescription and Patient Support Program Enrollment Form

Complete this form, which is available at www.adempas-us.com, and fax along with patient insurance information to the Adempas Program at 1-855-662-5200.

SECTION 1 Contact Information

By providing an email or phone, the patient agrees to receive automated calls, texts and/or email messages from myAIM Program administrator and the patient's pharmacy (CVS Specialty or Accredo) about your prescription and healthcare. Consent may be revoked at any time and is not a condition of the service. Carrier message and data rates may apply.

Patient Contact Information (* indicates required field)

Patient First Name*:	Patient Last Name*:			Birthdate* (MM/DD/YYYY):	Sex*: ☐ Male ☐ Female
Address*:	City*:	State*:	Zip Code*:	Preferred Phone*:	*OK to leave detailed message?* ☐ Yes ☐ No
Email: @	Preferred Language: ☐ English ☐ Spanish ☐ Other (specify) _____				
Alternate Contact Name:	Alternate Contact Phone:			Relationship to Patient:	

Prescriber Contact Information (* indicates required field)

Prescriber First Name*:	Prescriber Last Name*:			NPI*:
Address Line 1*:	Address Line 2:	City:	State:	Zip Code:
Office Contact:	Phone:	Fax:		

SECTION 2 Patient Information

Patient Information (* indicates required field)

Is Patient starting Adempas in a hospital setting? ☐ **Yes** ☐ **No** Start Date:_____ Discharge Date:_____

Does the patient have prescription coverage*? ☐ Yes ☐ No

Patient's local pharmacy:_____ Phone: _____

***PROVIDE ALL PATIENT INSURANCE INFORMATION, INCLUDING DRUG BENEFITS (FRONT AND BACK OF CARD) WITH THIS FORM.**

Please check one ICD-10 Code*:

Pulmonary arterial hypertension
- ☐ I27.0
- ☐ I27.21

Chronic thromboembolic pulmonary hypertension
- ☐ I27.24
  - ☐ Inoperable
  - ☐ Persistent/Recurrent

☐ **OTHER (please specify)**
_____
_____

Therapy Status:
- ☐ Initial therapy (monotherapy or in combination)
- ☐ Add-on therapy
- ☐ Transition from other therapy

Prescriber will comply with all Surescripts† terms and conditions including confidentiality, commercial messaging, privacy and security, applicable laws, and use of data. All Surescripts' disclaimers apply. A full list of terms and conditions is available at *https://ubc.com/surescriptsterms/*

SECTION 3 Prescription

Prescription (* indicates required field)

Note: NY Prescribers please submit prescription on an original NY State prescriptions blank. For all other States, send on a State-specific prescription blank if applicable for your State.

Starting dose*:	Titration schedule:
☐ **Adempas 1 mg tablet by mouth three times a day** ☐ Adempas 0.5 mg tablet by mouth three times a day Quantity: ☐ 30 day supply Refills:_____ Deliver to: ☐ Patient Home ☐ Prescriber Office	**Please check box for all dosages to be incorporated**: ☐ Based on patient's response per clinical evaluation of the physician or the nurse in consultation with the physician, the pharmacy is to provide the Adempas strength to accommodate titration needs of therapy. Adempas Tablets: 0.5 mg, 1 mg, 1.5 mg, 2 mg, 2.5 mg Directions: If systolic blood pressure is >95 mmHg and there are no signs/symptoms of hypotension, up titrate by 0.5 mg 3 times per day at intervals no sooner than 2 weeks to the highest tolerated dosage up to a maximum of 2.5 mg 3 times per day. If at any time, the patient has symptoms of hypotension, decrease the dosage by 0.5 mg 3 times daily. The established individual dose should be maintained. **Other special instructions:** _____ Titration Quantity: ☐ 30 day supply ☐ 14 day supply Refills: _____

I certify that the above information provided is accurate to the best of my knowledge. I appoint the Adempas AIM Program, on my behalf, to convey this prescription to the dispensing pharmacy. I understand that I may not delegate signature authority. Prescriber authorizes UBC to use the Surescripts Network† on Prescriber's behalf in connection with this prescription.

PRESCRIBER SIGNATURE REQUIRED		
	Dispense as Written*:	**Date*:**
	Substitutions Permitted*:	**Date*:**

Return this form along with patient insurance information to the Adempas Program via fax to 1-855-662-5200

†Surescripts is a consortium owned by some of the country's largest PBMs that offers information and technology services that support the electronic transmission of prescriptions between HCPs and other healthcare organizations.

To report any adverse events, product technical complaints, medication errors, or pregnancies associated with the use of Adempas, contact: Bayer at 1-888-842-2937, or send the information to DrugSafety.GPV.US@bayer.com.

INDICATIONS AND IMPORTANT SAFETY INFORMATION

INDICATIONS

- Adempas (riociguat) is indicated for the treatment of adults with persistent/recurrent chronic thromboembolic pulmonary hypertension (CTEPH) (WHO Group 4) after surgical treatment or inoperable CTEPH to improve exercise capacity and WHO functional class.
- Adempas is indicated for the treatment of adults with pulmonary arterial hypertension (PAH) (WHO Group 1) to improve exercise capacity, improve WHO functional class and to delay clinical worsening.*

Efficacy was shown in patients on Adempas monotherapy or in combination with endothelin receptor antagonists or prostanoids. Studies establishing effectiveness included predominantly patients with WHO functional class II–III and etiologies of idiopathic or heritable PAH (61%) or PAH associated with connective tissue diseases (25%).

*Time to clinical worsening was a combined endpoint defined as death (all-cause mortality), heart/lung transplantation, atrial septostomy, hospitalization due to persistent worsening of pulmonary hypertension, start of new PAH-specific treatment, persistent decrease in 6MWD, and persistent worsening of WHO functional class.

IMPORTANT SAFETY INFORMATION

WARNING: EMBRYO-FETAL TOXICITY

Adempas is contraindicated for use during pregnancy because it may cause fetal harm based on animal data.

Exclude pregnancy prior to the start of treatment with Adempas. Advise use of effective contraception prior to initiation of treatment, during treatment, and for one month after treatment with Adempas.

If pregnancy is detected, discontinue Adempas as soon as possible.

Contraindications

- Use of Adempas is contraindicated in patients who are pregnant.
- Co-administration of Adempas with nitrates or nitric oxide donors (such as amyl nitrite) in any form is contraindicated.
- Concomitant administration of Adempas with specific phosphodiesterase (PDE)-5 inhibitors (such as sildenafil, tadalafil, or vardenafil) or nonspecific PDE inhibitors (such as dipyridamole or theophylline) is contraindicated. Do not administer within 24 hours of sildenafil. Do not administer 24 hours before or within 48 hours after tadalafil.
- Adempas is contraindicated in patients with concomitant use of other soluble guanylate cyclase (sGC) stimulators.
- Adempas is contraindicated in patients with Pulmonary Hypertension associated with Idiopathic Interstitial Pneumonias (PH-IIP).

Warnings and Precautions

Embryo-Fetal Toxicity. Based on data from animal reproduction studies, Adempas may cause fetal harm when administered to a pregnant patient and is contraindicated during pregnancy. The available human data for riociguat do not establish the presence or absence of major birth defects related to the use of Adempas. Counsel patients who can become pregnant on the potential risk to a fetus. Exclude pregnancy prior to initiation of treatment with Adempas. Advise females of reproductive potential to use effective contraception prior to initiation of treatment, during treatment with Adempas and for at least one month after the last dose. When pregnancy is detected, discontinue Adempas as soon as possible.

Hypotension. Adempas reduces blood pressure. Consider the potential for symptomatic hypotension or ischemia in patients with hypovolemia, severe left ventricular outflow obstruction, resting hypotension, autonomic dysfunction, or concomitant treatment with antihypertensives or strong CYP and P-gp/BCRP inhibitors. Consider a dose reduction if patient develops signs or symptoms of hypotension.

Bleeding. In the placebo-controlled clinical trials, serious bleeding occurred in 2.4% of patients taking Adempas compared to 0% of placebo patients. Serious hemoptysis occurred in 5 (1%) patients taking Adempas compared to 0 placebo patients, including one event with fatal outcome. Serious hemorrhagic events also included 2 patients with vaginal hemorrhage, 2 with catheter-site hemorrhage, and 1 each with subdural hematoma, hematemesis, and intra-abdominal hemorrhage.

Pulmonary Veno-Occlusive Disease. Pulmonary vasodilators may significantly worsen the cardiovascular status of patients with pulmonary veno-occlusive disease (PVOD). Therefore, administration of Adempas to such patients is not recommended. Should signs of pulmonary edema occur, the possibility of associated PVOD should be considered and if confirmed, discontinue treatment with Adempas.

Most Common Adverse Reactions

The most common adverse reactions occurring more frequently (≥3%) on Adempas than placebo were headache (27% vs 18%), dyspepsia/gastritis (21% vs 8%), dizziness (20% vs 13%), nausea (14% vs 11%), diarrhea (12% vs 8%), hypotension (10% vs 4%), vomiting (10% vs 7%), anemia (7% vs 2%), gastroesophageal reflux disease (5% vs 2%), and constipation (5% vs 1%).

Other events that were seen more frequently in Adempas compared to placebo and potentially related to treatment were palpitations, nasal congestion, epistaxis, dysphagia, abdominal distension, and peripheral edema.

For important risk and use information, please click here to see the full Prescribing Information, including Boxed Warning.

To report any adverse events, product technical complaints, medication errors, or pregnancies associated with the use of Adempas, contact: Bayer at 1-888-842-2937, or send the information to DrugSafety.GPV.US@bayer.com.

PP-ADE-US-3398-1
August 2026

Phone: 1-855-4ADEMPAS (1-855-423-3672) Fax: 1-855-662-5200

Adempas® riociguat tablets
0.5mg | 1mg | 1.5mg | 2mg | 2.5mg

Adempas® (riociguat) Prescription and Patient Support Program Enrollment Form

Complete this form, which is available at www.adempas-us.com, and fax along with patient insurance information to the Adempas Program at 1-855-662-5200.

SECTION 4 Patient Support Program Enrollment

Patient Support Program Enrollment
Bayer offers patient support services for Adempas patients that include: (A) nurses to support you in starting therapy and achieving your optimal dose, (B) insurance benefit verification for Adempas and financial assistance for eligible patients and (C) education about CTEPH and/or PAH as well as helpful tips for managing your Adempas therapy ("myAIM"). These Programs are entirely optional and you may enroll in one or all of these Programs. To enroll in myAIM, you will need to sign a HIPAA authorization in order for your healthcare provider and/or pharmacy to share your protected healthcare information with Bayer and the myAIM Program administrator. You will remain enrolled in each Program that you select unless you opt-out by contacting myAIM via telephone at 1-855-423-3672 or until your HIPAA Authorization expires.
Please enroll me in: (check all that apply) ☐ A: Nursing ☐ B: Benefits Verification and Financial Assistance ☐ C: Digital Educational Information – e-mail address: _____@_____
Patient – please initial here to confirm your optional selections: _____ Patient can opt-out of any one of the above programs (or all) by contacting the AIM program.

THIS AREA INTENTIONALLY LEFT BLANK

To report any adverse events, product technical complaints, medication errors, or pregnancies associated with the use of Adempas, contact: Bayer at 1-888-842-2937, or send the information to DrugSafety.GPV.US@bayer.com.

PATIENT HIPAA AUTHORIZATION

I voluntarily provide this authorization for the use and disclosure of my Protected Health Information ("PHI"), as such term is defined by the Health Insurance Portability and Accountability Act of 1996 (as amended, "HIPAA"). I understand that PHI is health information that identifies me or that could reasonably be used to identify me.

I authorize my healthcare provider, including my physician and pharmacy, and my health plan, to disclose to Bayer and its contracted agents my name, address, telephone number, health insurance status and coverage and such medical information as may be necessary for me to enroll in the Aim Patient Support Program. I understand this disclosure(s) will contain PHI, including information about my current medical condition, treatment, coordination of treatment and receipt of medication. I allow the use and disclosure of my PHI to Bayer and its contracted agents for the following purposes:

- To verify my insurance information and coverage
- To ensure the accuracy and completeness of the Aim Patient Support Program Enrollment Form
- To help with my insurance coverage questions for Bayer medications
- To determine if I qualify for other Bayer patient support programs
- To determine my eligibility for other sources of prescription medication financial assistance
- To provide education, training, and ongoing support on the use of my Bayer medication
- To send me information on Bayer products and services related to my treatment
- To send me refill reminders for my Bayer prescription medication and to encourage its appropriate use
- To communicate with me, my healthcare providers and health plan about my medical care and treatment
- To contact me for market research feedback, sales support purposes, and as necessary to comply with applicable laws
- Bayer may contact me for potential adverse event follow-up information

I understand that:

- This Authorization will remain in effect until the end of my participation in the Aim Patient Support Program or 10 years, unless subject to applicable law from the date of my signature on this Authorization, whichever occurs later.
- I may cancel this Authorization at any time by contacting myAIM via telephone at 1-855-423-3672.
- If I cancel this Authorization my healthcare provider and health plan will stop sharing my PHI with Bayer and its contracted agents. However, the revocation will not affect prior use or disclosure of my PHI in reliance on this Authorization.
- I may opt-out of being contacted for market research feedback, sales support purposes, and still enroll in the patient support program.
- That entities that receive my PHI in accordance with this Authorization may not be required by law to keep the information private and that it will no longer be protected by the HIPAA privacy law. It may become available in the public domain.
- I do not need to sign this Authorization to receive (i) medical treatment or medication or (ii) coverage, payment, enrollment in or eligibility in benefits from my health plan. However, if I do not sign this Authorization, I may not participate in the Aim Patient Support Program or be eligible for other Bayer patient support programs.
- My healthcare providers, insurers, pharmacies and health plans may receive remuneration (payment) from Bayer in exchange for providing services to Bayer that may involve use or disclosure of my PHI.

I have read and understand the terms of this Authorization and have had the opportunity to ask questions about the uses and disclosures of PHI. I understand that I am entitled to receive a signed copy of this authorization and can get more information about the use and disclosure of PHI by contacting the Aim Patient Support Program at 1-855-4ADEMPAS (1-855-423-3672).

PATIENT TO SIGN AND DATE

Patient Name (print): _____

Patient (or legal guardian) Signature*: _____ **Date (mm/dd/yyyy):** _____

If signed by a legal representative —

Print Name: _____ **Relationship to patient:** _____

To report any adverse events, product technical complaints, medication errors, or pregnancies associated with the use of Adempas, contact: Bayer at 1-888-842-2937, or send the information to DrugSafety.GPV.US@bayer.com.